

Ateneo de Manila University

**Archium Ateneo**

---

Biology Faculty Publications

Biology Department

---

3-2016

## Challenges and Opportunities in the Culture of Giant Freshwater Prawn in the Philippines

Janice A. Ragaza

Follow this and additional works at: <https://archium.ateneo.edu/biology-faculty-pubs>



Part of the [Aquaculture and Fisheries Commons](#), and the [Biology Commons](#)

---

# CHALLENGES AND OPPORTUNITIES IN THE CULTURE OF GIANT FRESHWATER PRAWN IN THE PHILIPPINES

JANICE A. RAGAZA

Amid the irremediable decline of capture fisheries, the Philippine government has identified aquaculture as a sector that will ensure food security and generate new job opportunities geared toward the country's goal of economic development (Nagothu and Ortiz 2007). Aquaculture of giant freshwater prawn *Macrobrachium rosenbergii* (Fig. 1) is an emerging program that is intended to expand and vary the aquatic species used in freshwater aquaculture, which is still predominantly based on tilapia (Rosario and Tayamen 2004). Attractive characteristics of freshwater prawn culture include ease of breeding, high hatching rate, superior quality of meat and high market value (Keysami and Mohammadpour 2013).

Giant freshwater prawn is cultured in about 43 countries, with Asia representing more than 98 percent of global production (Mather and de Bruyn 2003). Principal producers are Bangladesh, China, India, Myanmar, Taiwan, Thailand and Vietnam (New and Nair 2012). It is widely distributed in most tropical and sub-tropical countries, including the Philippines. More than 100 species exist globally, 12 of which are found in the Philippines (Rosario and Tayamen 2004). The giant freshwater prawn is a dominant species, along with four others. It thrives in inland bodies of water in Agusan, Bicol region, Bulacan, Cagayan, Cotabato, Ilocos, Laguna, Lanao, Leyte, Maguindanao, some parts of Mindanao, Palawan,



FIGURE 1. Adult giant freshwater prawns at the Bureau of Fisheries and Aquatic Resources-National Freshwater Fisheries Technology Center, Muñoz City, Nueva Ecija, Philippines.



FIGURE 2. Experimental-scale culture of giant freshwater prawn in net cages at the Bureau of Fisheries and Aquatic Resources-National Freshwater Fisheries Technology Center, Muñoz City, Nueva Ecija Philippines.

Pampanga, Pangasinan, and Samar (Rosario and Tayamen 2004).

Names for freshwater prawn in local dialects are *ulang* (Bulacan, Laguna, Zambales and most parts of the country), *udang* (Ilocos, Cagayan and other parts of Northern Luzon), *urang* (Leyte), *budsang* (Bicol), *kalig* (Leyte), *kising-kising* (Pangasinan), *padao* (Cotabato), and *paje* (Palawan and Zamboanga) (Rosario and Tayamen 2004, Tayamen 2005).

The culture of freshwater prawn in the Philippines started in the 1970s. Culture trials ensued (Fig. 2), but technology was not sustained at the commercial level. Mass-production of seed and post-larvae were established by the Bureau of Fisheries and Aquatic Resources (BFAR) in the early 2000s. Since then, BFAR and other government agencies have conducted technology dispersal programs. Freshwater prawn production in ponds increased from 2.80 t in

2010 to 2.86 t in 2011 (BFAR 2010, 2011). The same trend was seen in aquaculture systems using small-farm reservoirs, which produced 1.56 t in the same year, an increase of 0.30 t from the previous year (BFAR 2010, 2011).

The giant freshwater prawn is considered to be the world's largest shrimp. In the wild, freshwater prawn can grow to 500

(CONTINUED ON PAGE 60)

g and fetch approximately US\$ 10/kg (Tayamen 2005). The freshwater prawn is gradually establishing a niche in local markets and also represents a multi-million dollar export opportunity to Japan, US, Taiwan and the European Union. Freshwater prawn is a high-value species that represents an alternative to black tiger shrimp *Penaeus monodon* and whiteleg shrimp *Litopenaeus vannamei*. Farmed freshwater prawns on average weigh from 30-100 g, which about the same size as medium to jumbo black tiger shrimp.

## INFECTIOUS DISEASES IN FRESHWATER PRAWN CULTURE

With the steady increase of global and local shrimp market prices in recent years, shrimp aquaculture has been converting from extensive and semi-intensive farming to super-intensive systems. This change in farming system can cause serious problems with infectious diseases because cultured shrimps are exposed to stressful conditions. Disease control and biosecurity are among the most critical issues that threaten the sustainability of commercial shrimp farming.

Giant freshwater prawn has been cultured for over 40 years, but there is scant information on pathogens and parasites (Arthur *et al.* 2005). Although it is a moderately disease-resistant species compared to penaeid shrimp (Ravi *et al.* 2009), with the rapid development of intensive farming, microorganisms have been implicated in many serious diseases (Liang *et al.* 2011).

The common causes of infectious diseases in giant freshwater prawn are mostly viruses and bacteria (*Vibrio* and *Aeromonas*) that impede production of seed and cause stock mortalities (Bachere 2000). Early larval stages are very susceptible to vibriosis and *Aeromonas* infections, which can cause complete mortality (Tonguthai 1997). The production and survival of healthy larvae and post-larvae has been a primary hindrance in the expansion of freshwater prawn aquaculture (Prakash and Karmagam 2013).

## USE OF IMMUNOSTIMULANTS IN SHRIMP AQUACULTURE

The crustacean immune system lacks adaptive immune response mechanisms and therefore defense depends on the non-specific immune response. The non-specific immune defense of crustaceans is primarily attributed to hemocytes, which are specialized blood cells that carry out cellular and humoral mechanisms such as phagocytosis and encapsulation, and production of antimicrobial substances to combat foreign particles and infectious agents, among others (Lio-Pio and Inui 2010). Additionally, crustaceans upregulate expression of immune-related genes in response to viruses and other pathogens.

One promising technique for prophylaxis and disease control is to increase the immune status of shrimp with immunostimulants. According to Bricknell and Dalmo (2005), "An immunostimulant is a naturally occurring compound that modulates the immune system by increasing the host's resistance against diseases that in most circumstances are caused by pathogens." This approach enhances innate immune responses, resistance to bacterial and viral infections, resistance to pathological and environmental stresses, and growth performance in many shrimp species (Wang *et al.* 2013).

Immunostimulants used in aquaculture provoke beneficial

effects because of enhanced cellular and humoral responses in cultured species (Ringø *et al.* 2012). Immunostimulants influence and target non-specific immune aspects such as complement levels, lysozyme levels, natural killer activity, phagocytic cell activity, and total immunoglobulin levels, among others (Shankar *et al.* 2012).

Different immunostimulants activate various aspects of the immune system and their efficacies are species-specific, pathogen-specific and largely depend on delivery method (Sakai 1999, Smith *et al.* 2003, Agrawal *et al.* 2010). Immunostimulants are administered by immersion, injection or through dietary inclusion (Rocha-Montero *et al.* 2006, Chiu *et al.* 2010, Wang *et al.* 2013). Dietary immunostimulants offer a cheap and effective method for increasing the productivity and cost-effectiveness of aquaculture operations by enhancing disease resistance and stress tolerance and reducing the need for more radical and costly disease control measures.

A number of substances confer immunostimulatory properties in different species of cultured shrimp. Immunostimulants include inactivated microbes and their cellular components, complex carbohydrates and polysaccharides, animal extracts, cytokines, lectins, plant and herbal extracts, vitamins, minerals and synthetic compounds.

## OUTLOOK FOR THE FUTURE

Factors that constrain development of giant freshwater prawn culture in the Philippines include inadequate information on nutritional requirements of the prawn, the need for broodstock and seed improvement, lack of technological innovations that lower production cost and improve its growth and survival, and low extension of technology to potential farmers (Rosario and Tayamen 2004, Tayamen 2005, De Guzman 2008). Freshwater prawn aquaculture continues to face serious risks and threats of diseases, which explains in part the decrease in global production of prawns (Angeles *et al.* 2009, Kutty 2005).

The risk and threat of disease outbreaks can be reduced with the use of dietary immunostimulants. This practice can support good production levels of freshwater prawn when disease resistance through dietary stimulation is achieved. Development of this industry can certainly translate to employment opportunities, providing prawn farmers with stable jobs and livelihoods.

## Notes

Janice A. Ragaza, Department of Biology, School of Science and Engineering, Loyola Schools, Ateneo de Manila University, Katipunan Avenue, Loyola Heights, Quezon City 1108, Manila, Philippines. E-mail: jragaza@ateneo.edu

## References

- Agrawal, S., S. Khadase and G. Talele. 2010. Bioactive immunomodulatory fraction from *Tridaxprocumbens*. *Asian Journal of Biological Sciences* 3:120-127.
- Angeles, I., Y.H. Chien and M.M. Tayamen. 2009. Effects of different dosages of astaxanthin on giant freshwater prawn *Macrobrachium rosenbergii* (De Man) challenged with *Lactococcus garvieae*. *Aquaculture Research* 41:70-77.
- Arthur, J.R., D. Hurwood, E.R. Lovell, M.G. Bondad-Reantaso and P.B. Mather. 2005. Pathogen and ecological risk analysis for the

- introduction of giant river prawn, *Macrobrachium rosenbergii*, from Fiji to the Cook Islands. Aquaculture Technical Paper, Secretariat of the Pacific Community.
- Bachere, E. 2000. Shrimp immunity and disease control. *Aquaculture* 191:3-11.
- BFAR (Bureau of Fisheries and Aquatic Resources). 2010. Philippine fisheries profile 2010. Department of Agriculture, Bureau of Fisheries and Aquatic Resources, Diliman, Quezon City, Philippines.
- BFAR (Bureau of Fisheries and Aquatic Resources). 2011. Philippine Fisheries Profile 2011. Department of Agriculture, Bureau of Fisheries and Aquatic Resources, Diliman, Quezon City, Philippines.
- Bricknell, I. and R.A. Dalmo. 2005. The use of immunostimulants in fish larval aquaculture. *Fish and Shellfish Immunology* 19:457-472.
- Chiu, S.T., S.L. Hsieh, S.P. Yeh, S.J. Jian, W. Cheng and C.H. Liu. 2010. The increase of immunity and disease resistance of the giant freshwater prawn, *Macrobrachium rosenbergii* by feeding with selenium enriched-diet. *Fish and Shellfish Immunology* 29:623-629.
- De Guzman, C.B. 2008. A viable alternative to ulang pond culture. Bureau of Agricultural Research (BAR) Digest 10:3.
- Keysami, M.A. and M. Mohammadpour. 2013. Effect of *Bacillus subtilis* on *Aeromonas hydrophila* infection resistance in juvenile freshwater prawn, *Macrobrachium rosenbergii* (de Man). *Aquaculture International* 21:553-562.
- Kutty, M.N. 2005. Towards sustainable freshwater prawn aquaculture — lessons from shrimp farming, with special reference to India. *Aquaculture Research* 36:255-263.
- Liang, T., X. Li, J. Du, W. Yao, G. Sun, X. Dong, Z. Liu, J. Ou, Q. Meng, W. Gu and W. Wang. 2011. Identification and isolation of a spiroplasma pathogen from diseased freshwater prawns, *Macrobrachium rosenbergii*, in China: a new freshwater crustacean host. *Aquaculture* 318:1-6.
- Lio-Pio, G.D. and Y. Inui. 2010. Health Management in Aquaculture, 2nd ed. Southeast Asian Fisheries Development Center/Aquaculture Department (SEAFDEC/AQD), Tigbauan, Iloilo.
- Mather, P.B. and M. de Bruyn. 2003. Genetic diversity in wild stocks of the giant freshwater prawn (*Macrobrachium rosenbergii*): implications for aquaculture and conservation. *World Fish Center Quarterly* 26(4):4-7.
- Nagothu, U.S. and I. Ortiz. 2007. Review and assessment of socio-economic issues of aquaculture in the Philippines: mitigating impact of aquaculture in the Philippines. Norwegian Institute for Water Research and PHILMINAQ.
- New, M.B. 2002. Farming freshwater prawns. A manual for the culture of giant river prawn (*Macrobrachium rosenbergii*). FAO Fisheries Technical Paper 428. Rome, FAO.
- New, M.B. and C.M. Nair. 2012. Global scale of freshwater prawn farming. *Aquaculture Research* 43:960-969.
- Prakash, M. and N. Karmagam. 2013. A study on bacterial flora associated with freshwater prawn, *Macrobrachium rosenbergii*. *International Journal of Current Research and Academic Review* 1:1-16.
- Ravi, M., A. Nazeer Basha, M. Sarathi, H.H. Rosa Idalia, J. Sri Widada and J.R. Bonami. 2009. Studies on the occurrence of white tail disease (WTD) caused by MrNV and XSV in hatchery-reared post-larvae of *Penaeus indicus* and *P. monodon*. *Aquaculture* 292:117-120.
- Ringø, E., R.E. Olsen, J.L.G. Vecino, S. Wadsworth and S.K. Song. 2012. Use of immunostimulants and nucleotides in aquaculture: a review. *Journal of Marine Science Research and Development* 2:1.
- Rocha-Montero, A., D. McIntosh, R. Sanchez-Merino and I. Flores. 2006. Immunostimulation of white shrimp (*Litopenaeus vannamei*) following dietary administration of Ergosan. *Journal of Invertebrate Pathology* 91:188-194.
- Rosario, W.R. and M.M. Tayamen. 2004. Culture of giant freshwater prawn: Philippines. Southeast Asian Fisheries Development Center/Aquaculture Department (SEAFDEC/AQD) Institutional Repository (SAIR).
- Sakai, M. 1999. Current research status of fish immunostimulants. *Aquaculture* 172:63-92.
- Shankar, R., H. Shivananda Murthy, H.R. Sujatha, E.G. Jayaraj, C.S. Tejpal and V.S. Chinthamani. 2012. Effect of nucleotide on growth, immune responses and resistance of *Macrobrachium rosenbergii* (De Man) to *Macrobrachium rosenbergii* nodavirus (MrNV) and extra small virus (XSV) and *Aeromonas hydrophila* infection. *Aquaculture International* 20:1-12.
- Smith, V.J., J.H. Brown and C. Hauton. 2003. Immunostimulation in crustaceans: Does it really protect against infection? *Fish and Shellfish Immunology* 15:71-90.
- Tayamen, M.M. 2005. Freshwater prawn program of Bureau of Fisheries and Aquatic Resources (BFAR). Southeast Asian Fisheries Development Center/Aquaculture Department (SEAFDEC/AQD) Institutional Repository (SAIR).
- Tonguthai, K. 1997. Diseases of the freshwater prawn, *Macrobrachium rosenbergii*. The Aquatic Animal Health Research Institute 4(2).
- Vici, V., B. Singh and S.G. Bhat. 2000. Application of bacterins and yeast *Acremonium dyosporii* to protect the larvae of *Macrobrachium rosenbergii* from vibriosis. *Fish and Shellfish Immunology* 10:559-563.
- Wang, Y.C., C.F. Chang and H.Y. Chen. 2013. The role of glucans in protection of shrimp against disease. In V. Vetvicka and M. Novac, editors. *Biology and Chemistry of Beta-glucan*. Biology and Chemistry of Beta-glucan 2:173-195.

THE RISK AND THREAT OF DISEASE OUTBREAKS CAN BE REDUCED WITH THE USE OF DIETARY IMMUNOSTIMULANTS. THIS PRACTICE CAN SUPPORT GOOD PRODUCTION LEVELS OF FRESHWATER PRAWN WHEN DISEASE RESISTANCE THROUGH DIETARY STIMULATION IS ACHIEVED. DEVELOPMENT OF THIS INDUSTRY CAN CERTAINLY TRANSLATE TO EMPLOYMENT OPPORTUNITIES, PROVIDING PRAWN FARMERS WITH STABLE JOBS AND LIVELIHOODS.